



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ANSWER TO QUERY AT PAGE 143, VOL. V.

BY GEORGE EASTWOOD, SAXONVILLE, MASS.

Analysis.—Suppose the construction effected, and that the required circles O , O' , O'' , are drawn so as to touch each other and the given circles Q and Q' , as in the annexed diagram.

Join the centers O and O'' ; O' and O'' ; O and Q' ; and O' and Q' . Produce OO'' to L , and OQ' to H , and upon AD demit the perpendiculars $O''K$ and $Q'M$.

By a property of the Arbelos (See Leybourn's Math. Rep., pp. 207–209), $O''K = 2FO''$, $Q'M = 4EQ'$, etc. [See ANALYST, p. 171, Vol. I.]

From the triangle $O'O''O$, we have

$$\begin{aligned} (OF + FO'')^2 &= O'O^2 + O''O^2 + 2O'O.OK \\ &= (AO - AO')^2 + (OL - O'L)^2 + 2(AO - AO')OK \\ &= (AO - AO')^2 + (AO - FO'')^2 + 2(AO - AO')OK; \end{aligned}$$

whence
$$OK = \frac{(AO + AO')FO''}{AO - AO'} - AO.$$

Make $AO = R$, $AO' = R'$, $BQ = R''$, $FO'' = r$, and $EQ' = r'$; then, since $R = R' + R''$, we have

$$R + R' = 2R' + R'', \quad R - R' = R'',$$

and therefore
$$OK = \frac{(2R' + R'')r}{R'} - (R' + R''). \quad (A)$$

But $OK^2 = OO''^2 - O''K^2 = OO''^2 - (2FO'')^2 = (R' + R'' - r)^2 - 4r^2. \quad (B)$

Putting the square of (A) in (B), we deduce


$$r = \frac{(R' + R'')R'R''}{R'^2 + (R' + R'')R'}. \quad (C)$$

By a similar process we deduce

$$r' = \frac{(R' + R'')R'R''}{4R'^2 + (R' + R'')R'}. \quad (D)$$

By the question, $r' = EQ' = Q'H$, is a given line. Therefore, solving the symbolical equation (D) for R' , we readily find

$$\begin{aligned} R' &= \frac{1}{2}R''[\sqrt{(R'' + 15r') - 1}], \\ R &= \frac{1}{2}R''[\sqrt{(R'' + 15r') + 1}], \\ r &= \frac{R''(R'' + 15r' - 1)}{R'' + 15r' + 3}. \end{aligned}$$



Furthermore, adding AO to OK we find

$$AK = \frac{(AO + AO')FO''}{AO - AO'} = \frac{(2R' + R'')r}{R''} = \frac{R''(R'' + 15r')^{\frac{1}{2}}(R'' + 15r' - 1)}{R'' + 15r' + 3},$$

and, similarly,

$$AM = \frac{(AO + AO')EQ'}{AO - AO'} = \frac{(2R' + R'')r'}{R''} = r'\sqrt{(R'' + 15r')}. \quad \text{Hence this}$$

Construction.—On the indefinite straight line AX , set off

$$AB = 2AO' = \frac{1}{2}R''[\sqrt{(R'' + 15r')} - 1],$$

$$AD = 2AO = \frac{1}{2}R''[\sqrt{(R'' + 15r')} + 1],$$

$$BQ = R''.$$

Bisect AB in O' , and AD in O : then, with centers O' , O and Q , and with radii AO' , AO and BQ , describe the circles AEB , $AHLD$ and BGD . These will obviously touch each other, two and two.

Again, on AD set off

$$AK = \frac{R''\sqrt{(R'' + 15r')(R'' + 15r' - 1)}}{R'' + 15r' + 3},$$

and

$$\begin{aligned} AM &= r'\sqrt{(R'' + 15r')}. \quad \text{At } K, \text{ erect the perpend. } KO'' \\ &= \frac{2R''(R'' + 15r' - 1)}{R'' + 15r' + 3}, \end{aligned}$$

and at M erect the perpendicular $MQ' = 4r'$. Finally, with centers O'' and Q' , and radii $FO'' = R''(R'' + 15r' - 1) \div (R'' + 15r' + 3)$ and $EQ' = 4r'$, describe the circles FIL , EIH and it is done.

[The foregoing solution, it will be seen, applies only to a special case of the question quoted at p. 143, Vol. V, as the relative position of the given circles cannot be arbitrarily taken. No general solution has been rec'd.]

NOTE ON THE DIFFERENTIATION OF EXPONENTIAL AND TRIGONOMETRICAL FUNCTIONS.

BY ARTEMAS MARTIN, M. A., ERIE, PA.

THE following methods may not be new, but I have not met with them in any of the many works on the calculus to which I have access.

1. Differentiate $y = a^x$.

We have, by the Exponential Theorem,

$$y = a^x = 1 + x \log a + \frac{x^2(\log a)^2}{1.2} + \frac{x^3(\log a)^3}{1.2.3} + \text{etc};$$